

# THE ULTRAVIOLET FLASH ACCOMPANYING GRBS FROM NEUTRON-RICH INTERNAL SHOCKS

Y. Z. FAN<sup>1,2,3</sup> AND D. M. WEI<sup>1,2</sup>

<sup>1</sup>Purple Mountain Observatory, Chinese Academy of Science, Nanjing 210008, China

<sup>2</sup>National Astronomical Observatories, Chinese Academy of Sciences, Beijing, 100012, China

<sup>3</sup>On leave at Dept. of Physics, University of Nevada, Las Vegas, NV89154, USA

yzfan@pmo.ac.cn (yzfan@physics.unlv.edu) dmwei@pmo.ac.cn

*accepted for publication in ApJL*

## ABSTRACT

In the neutron-rich internal shocks model for Gamma-ray Bursts (GRBs), the Lorentz factors (LFs) of ion shells are variable, so are the LFs of accompanying neutron shells. For slow neutron shells with a typical LF  $\sim$  tens, the typical  $\beta$ -decay radius reads  $R_{\beta,s} \sim \text{several} \times 10^{14} \text{cm}$ , which is much larger than the typical internal shocks radius  $\sim 10^{13} \text{cm}$ , so their impact on the internal shocks may be unimportant. However, as GRBs last long enough ( $T_{90} > 20(1+z)\text{s}$ ), one earlier but slower ejected neutron shell will be swept successively by later ejected ion shells in the range  $\sim 10^{13} \text{cm} - 10^{15} \text{cm}$ , where slow neutrons have decayed significantly. We show in this work that ion shells interacting with the  $\beta$ -decay products of slow neutron shells can power a ultraviolet (UV) flash bright to 12th magnitude during the prompt  $\gamma$ -ray emission phase or slightly delayed, which can be detected by the upcoming Satellite SWIFT in the near future.

*Subject headings:* gamma rays:bursts — radiation mechanisms: nonthermal — shock waves

## 1. INTRODUCTION

As realized by many authors, the fireball of Gamma-ray bursts (GRBs) may contain significant neutron component in essentially all progenitor scenarios (e.g., Derishev, Kocharovsky & Kocharovsky 1999a, b; Beloborodov 2003a). The dynamics as well as the observable signatures (the neutrino emission from the proton-neutron decoupling and the changes in GRB lightcurves due to neutrons preceding the ion shell or catching up it) of a relativistic neutron-rich fireball have been investigated by Derishev, Kocharovsky & Kocharovsky (1999a, b) firstly. Since then, many authors have had their attention on the neutron-fed GRBs. If the neutron abundance is comparable to that of proton, the inelastic collision between differentially streaming protons and neutrons in the fireball will provide us observable 5-10 GeV neutrinos (Bachall & Mészáros 2000, hereafter BM00; Mészáros & Rees 2000). The baryon-loading problem in GRB models can be ameliorated if significant fraction of baryons confined in the fireball are converted to neutrons (Fuller, Pruet & Abazajian 2000). Further investigations on the observation implications have been presented in Pruet & Dalal (2002, hereafter PD02) and Beloborodov (2003b, hereafter B03). In PD02, the short GRBs are assumed to be powered by external shocks and the neutron component lags behind the ion one. As the ion ejecta has been decelerated significantly by the external medium, the decayed neutron ejecta catches up the decelerated ion ejecta and powers keV emission. In B03, the neutron-ejecta keeps ahead the ion-ejecta. The  $\beta$ -decay ( $n \rightarrow p + e^- + \bar{\nu}_e$ ) products share their momentum with the external medium immediately. As a result, the external medium have been accelerated to a ultra-relativistic velocity. The interaction between the ion ejecta and the accelerated medium is much different from the usual case (i.e., a fireball ejecta interacting with the static medium). So the presence of neutron ejecta qual-

itatively changes the early dynamical evolution of GRBs' remnant.

In PD02 and B03, only the “fast” neutron component has been taken into account. In fact, the long, complex GRBs are more likely to be powered by the interaction of shells with variant LFs, i.e., the internal shocks model (Paczynski & Xu 1994; Rees & Mészáros 1994). The best fit to the multi-wavelength afterglows implies that a significant fraction of the initial kinetic energy has been converted into internal energy (Panaitescu & Kumar 2002), which requires that the difference in velocities between the shells is significant (i.e., the corresponding LFs satisfy  $\eta_f \gg \eta_s$ ) and their masses are comparable ( $m_f \approx m_s$ ) (Piran 1999, hereafter P99) [Throughout this work, the subscript f, s represent the fast/slow shells respectively; n, p represent the neutron/proton component respectively]. Typically,  $\eta_s$  is in order of tens, and  $\eta_f$  is in order of hundreds. Thus, for the slow shell, the n, p component coast with  $\Gamma_{n,s} = \Gamma_{p,s} \simeq \eta_s$  since  $\eta_s < \eta_\pi \simeq 3.9 \times 10^2 L_{52}^{1/4} r_{0,7}^{-1/4} [(1+\xi)/2]^{-1/4}$  (BM00), where  $L$  being the total luminosity of the ejecta,  $r_0$  being the radius of the central engine,  $\xi$  being the ratio of the neutrons to the protons contained in the shells. In this Letter, we adopt the convention  $Q_x = Q/10^x$  for expressing the physical parameters, using cgs units. For the fast shell, the n, p components move with different LFs:  $\Gamma_{n,f} < \Gamma_{p,f}$  since generally  $\eta_f > \eta_\pi$  (BM00). In the internal shocks phase, the fast ion shell catches the slower but earlier one at a radius  $R_{\text{int}} \sim 10^{13} \text{cm}$  and forms a new one moving with a LF  $\Gamma_m \sim \text{several hundred}$  (hereafter the formed “new” shell is named as the “i-shell”). At  $R_{\text{int}}$ , the  $\beta$ -decay of the neutron component is far from important unless the typical variability timescale is longer than 0.1 s (See Rossi, Beloborodov & Rees 2004 for more information).

As the duration of GRB is long enough, the much later ejected i-shell catches up the earlier slow neutron shell

(hereafter, the “n-shell”) at  $R_{\text{cat}} \approx 2\Gamma_{\text{n,s}}^2 c\delta T/(1+z)$ , where  $\delta T$  being the ejection time-lag of the earlier slow n-shell and the later i-shell; On the other hand, the  $\beta$ -decay radius of slow neutrons reads  $R_{\beta,s} \approx 2.6 \times 10^{13} \Gamma_{\text{n,s}} \text{ cm}$ , at which slow neutrons have decayed significantly. As long as  $R_{\text{cat}}$  is comparable with  $R_{\beta,s}$ , i.e.,  $\delta T \geq 14(1+z)\Gamma_{\text{n,s},1.5}^{-1}$ , the decayed products of earlier but slower ejected neutron shells will be swept orderly by the later ejected ion shells in a range  $\simeq R_{\text{int}} - R_{\beta,s}$ . The possible emission signature powered by that interaction is of our interest in this Letter. Otherwise, for GRBs much shorter than  $14(1+z)\Gamma_{\text{n,s},1.5}^{-1}$ , the last i-shell crosses the first slow n-shell at a radius  $\ll R_{\beta,s}$ , where the  $\beta$ -decay is unimportant and there is no observable signature.

## 2. THE SIMPLEST MODEL

Even for a neutron-free internal shocks model, the practical LF distribution of shells involved is far from clear so far, let alone say the neutron-rich one. Here we use an approximate model proposed by Guetta, Spada & Waxman (2001), in which the LFs of i-shells are drawn from a bimodal distribution,  $\Gamma_{\text{ej}} = \eta_{\text{f}}$  or  $\Gamma_{\text{ej}} = \eta_{\text{s}}$ , with equal probability. That simple model is favored by its relative high energy conversion efficiency and peak energy. In this Letter, mainly for simplicity, we make the following assumptions: (i)  $\xi = 1$ , and the ratios between the mass of the fast neutrons, slow neutrons and i-shells are 1:1:2. (ii) Each fast (slow) neutron/ion shell moves with a same LF  $\Gamma_{\text{n,f}} \sim 300$  ( $\Gamma_{\text{n,s}} \sim 30$ ) and  $\Gamma_{\text{p,f}} \sim 1000$  ( $\Gamma_{\text{p,s}} \sim 30$ ) respectively. After the merger of a pair of fast/slow ion shells, the resulting i-shell moves with  $\Gamma_{\text{m}} \sim 200$ . (iii) The energetic ejecta are expanding into the low density interstellar medium (ISM), which is the case for most GRBs (Panaitescu & Kumar 2002). (iv) The duration of the GRB is longer than  $30(1+z)$ s since the constraint  $\delta T \simeq 14(1+z)\Gamma_{\text{n,s},1.5}^{-1}$  s should be satisfied.

The  $\beta$ -decay radius of fast neutrons  $R_{\beta,f} \approx 8 \times 10^{15} \Gamma_{\text{n,f},2.5} \text{ cm}$  is much larger than  $R_{\beta,s}$ . We ignore them since for  $R \leq R_{\beta,s}$ , there is only a small fraction of fast neutrons decayed. The ISM has been swept by the early and fast ions as well as the decayed products of fast neutrons, so the decay products of slow neutrons move freely.

Assuming that at radius  $R$ , the  $(k-1)$ th i-shell crosses the  $j$ th slow n-shell ( $j, k \gg 1$ ). At radius  $\simeq R + \Delta R$ , the  $k$ th i-shell will catch up the  $j$ th slow n-shell, where  $\Delta R \approx 2\Gamma_{\text{n,s}}^2 \Delta$ ,  $\Delta \approx c\delta t/(1+z)$ ,  $\delta t \sim 0.01(1+z)$ s being the typical variability timescale of GRBs. For the  $j$ th n-shell, the products of slow neutrons decayed at radius  $\leq R$  have been carried away by i-shells ejected early than the  $k$ th one. Hence the mass swept by  $k$ th i-shell is  $\Delta M \simeq \frac{M_{\text{n}} \Delta R}{R_{\beta,s}}$ , where  $M_{\text{n}} \equiv M_{\text{n}}^0 \exp(-\frac{R}{R_{\beta,s}})$ ,  $M_{\text{n}}^0$  being the the initial rest mass of the n-shell. So the averaged comoving number density of proton (due to  $\beta$ -decay) swept by  $k$ th i-shell can be estimated by

$$n \simeq \frac{\Delta M}{4\pi R^2 m_{\text{p}} \Gamma_{\text{n,s}} \Delta} \approx \frac{\Gamma_{\text{n,s}} M_{\text{n}}}{2\pi R^2 m_{\text{p}} R_{\beta,s}}. \quad (1)$$

Notice that in front of the much later ejected i-shells, there are hundreds of decaying n-shells. With assumptions made before, the whole process of each i-shell interacting with these decaying n-shells is rather similar. For convenience, in our following treatment, the discrete interaction

of each i-shell with these decaying n-shells has been simplified as an i-shell sweeping a moving proton trail (the LF of which is  $\Gamma_{\text{n,s}}$ ) with a number density  $n$  continually. The dynamics of that interaction can be described by the shock model as follows: Now the trail is moving with a LF  $\Gamma_{\text{n,s}} \sim \text{tens}$ , the generated thermal energy (comoving frame) in the shock front satisfies  $dU = (\gamma_{\text{rel}} - 1)dm c^2$  (e.g., B03) rather than  $dU = (\gamma - 1)dm c^2$ ; where  $\gamma_{\text{rel}} = \gamma\Gamma_{\text{n,s}}(1 - \beta_{\gamma}\beta_{\Gamma_{\text{n,s}}})$  being the LF of the decelerating i-shell (the LF of which is  $\gamma$ ) relative to the trail;  $\beta_{\gamma}$  and  $\beta_{\Gamma_{\text{n,s}}}$  being the corresponding velocity of  $\gamma$  and  $\Gamma_{\text{n,s}}$ ; the mass swept by the i-shell reads

$$dm = 4\pi n m_{\text{p}} \Gamma_{\text{n,s}} R^2 (\beta_{\gamma} - \beta_{\Gamma_{\text{n,s}}}) dR \simeq \frac{M_{\text{n}}}{R_{\beta,s}} dR. \quad (2)$$

After some simple algebra, the energy conservation of the system (the decelerating i-shell and the swept slow neutron trail) yields

$$\frac{d\gamma}{dm} = -\frac{\gamma\gamma_{\text{rel}} - \Gamma_{\text{n,s}}}{M_{\text{ion}} + m + (1 - \epsilon)U}, \quad (3)$$

where  $M_{\text{ion}}$  being the rest mass of the i-shell;  $\epsilon$  being the radiation efficiency of the shock. For  $\Gamma_{\text{n,s}} = 1$ , equation (3) reduces to the familiar form  $\frac{d\gamma}{dm} = -\frac{\gamma^2 - 1}{M_{\text{ion}} + m + (1 - \epsilon)U}$  (P99).

In the downstream, the electrons have been heated by the shock. As usual (e.g., P99), we assume that the shocked electrons distribute as  $\frac{dn}{d\gamma_e} \propto \gamma_e^{-p'}$  for  $\gamma_e > \gamma_{e,m}$ , where  $p' \sim 2.3$  being the typical power law index,  $\gamma_{e,m} \approx \epsilon_e \frac{p'-2}{p'-1} \frac{m_{\text{p}}}{m_e} (\gamma_{\text{rel}} - 1)$ ;  $\epsilon_e$  and  $\epsilon_B$  being the fractions of the shock energy given to electrons and the random magnetic field at shock, respectively. The comoving downstream magnetic field  $B \approx \sqrt{32\pi\epsilon_B\gamma_{\text{rel}}(\gamma_{\text{rel}} - 1)n m_{\text{p}} c^2}$ . As shown in P99, there is a critical LF, above which synchrotron radiation is significant:  $\gamma_{e,c} = \frac{6\pi m_e c}{(1+Y)\sigma_{\text{T}}\gamma B^2 t}$ , where  $t = t_{\text{obs}}/(1+z)$  is determined by  $dR = 2\gamma^2 c dt$  ( $t_{\text{obs}}$  being the observer time,  $z \sim 1$  being the red-shift of GRBs),  $\sigma_{\text{T}}$  being the Thomson cross section,  $Y$  being the Compton parameter (e.g., Wei & Lu 1998, 2000; Sari & Esin 2001):  $Y \simeq \frac{-1 + \sqrt{1 + 4x\epsilon_e/\epsilon_B}}{2}$ , where  $x$  being the radiation coefficient of electrons, so  $\epsilon \equiv x\epsilon_e$ . For  $\gamma_{e,m} > \gamma_{e,c}$  (fast cooling),  $x = 1$ ; For  $\gamma_{e,m} < \gamma_{e,c}$ ,  $x = (\gamma_{e,m}/\gamma_{e,c})^{\frac{p'-2}{2}}$  (slow cooling).

Given a proper boundary condition, equation (3) can be easily solved. With resulting  $\gamma_{e,m}$ ,  $\gamma_{e,c}$  and  $B$ , we can calculate the typical synchrotron radiation frequency  $\nu_{\text{m}} = \frac{\gamma_{e,m}^2 \gamma_e B}{2(1+z)\pi m_e c}$  ( $e$  being the charge of electron), the cooling frequency  $\nu_c = \frac{\gamma_{e,c}^2 \gamma_e B}{2(1+z)\pi m_e c}$  and the self-absorption frequency  $\nu_a$  (the detailed calculation on  $\nu_a$  can be found in the appendix of Wu et al. (2003)), with which we can analytically calculate the synchrotron radiation flux at the fixed band. The numerical results for these typical frequencies have been plotted in figure 1, where the parameters are taken as:  $M_{\text{ion}} = 2M_{\text{n}}^0 = 5.6 \times 10^{26} \text{ g}$ , which is corresponding to a Luminosity  $L = 10^{52} \text{ ergs s}^{-1}$  and  $\delta t = 10^{-2}(1+z)$ s,  $\Gamma_{\text{m}} = 200$ ;  $\Gamma_{\text{n,s}} = 30$ ,  $\epsilon_e = 0.3$ ,  $\epsilon_B = 0.01$ ,  $z = 1$  ( $D_{\text{L}} = 2.2 \times 10^{28} \text{ cm}$ ). As shown in figure

1, at the early time, the electrons are usually in fast cooling case and  $\nu_a$  is above the optical band. Lately, the electrons are in slow cooling case and  $\nu_a$  drops to  $\sim$  a few  $\times 10^{13}$  Hz (Currently, the observed emission are coming from hundreds of i-shells interacting with the slow neutrons trail. For the same reason, in our estimating the self-absorption frequency, the total number of electrons contributed has been assumed to be 300 times that of one trail. In fact, if just an i-shell interacting with the slow neutron trail has been taken into account,  $\nu_a$  should be one order or more less than the value presented here).

The synchrotron flux as a function of observer frequency can be approximated as follows: In the case of fast cooling (Just for  $\nu_a > \min\{\nu_c, \nu_m\}$ . For  $\nu_a < \min\{\nu_c, \nu_m\}$ , please refer to Eqs. (4-5) of Zhang & Mészáros (2004)),

$$F_\nu \approx F_{\max} \begin{cases} (\nu_c/\nu_a)^3(\nu/\nu_c)^2, & (\nu < \nu_c), \\ (\nu_a/\nu_c)^{-1/2}(\nu/\nu_a)^{5/2}, & (\nu_c < \nu < \nu_a), \\ (\nu/\nu_c)^{-1/2}, & (\nu_a < \nu < \nu_m), \\ (\nu_m/\nu_c)^{-1/2}(\nu/\nu_m)^{-p'/2}, & (\nu > \nu_m). \end{cases} \quad (4)$$

In the case of slow cooling

$$F_\nu \approx F_{\max} \begin{cases} (\frac{\nu_m}{\nu_a})^{\frac{(p'+4)}{2}}(\frac{\nu}{\nu_m})^2, & (\nu < \nu_m), \\ (\frac{\nu_a}{\nu_m})^{-\frac{(p'-1)}{2}}(\frac{\nu}{\nu_a})^{5/2}, & (\nu_m < \nu < \nu_a), \\ (\frac{\nu}{\nu_m})^{-\frac{(p'-1)}{2}}, & (\nu_a < \nu < \nu_c), \\ (\frac{\nu_c}{\nu_m})^{-\frac{(p'-1)}{2}}(\frac{\nu}{\nu_c})^{-\frac{p'}{2}}, & (\nu > \nu_c). \end{cases} \quad (5)$$

Approximately,  $F_{\max} = \frac{3\sqrt{3}\Phi_{p'}(1+z)N_e m_e c^2 \sigma_T}{32\pi^2 e D_L^2} \gamma B$ ,  $\Phi_{p'}$  being a function of  $p'$  (for  $p' \sim 2.3$ ,  $\Phi_{p'} \simeq 0.6$ );  $N_e$  being the number of electrons involved in the emission,  $D_L$  being the luminosity distance (Wijers & Galama 1999).

The upcoming Satellite Swift carries three telescopes<sup>1</sup>: The Burst Alert Telescope (BAT), the X-ray Telescope (XRT), the Ultraviolet and Optical Telescope (UVOT). The energy range of XRT is 0.2-10 keV. UVOT covers 170nm-650nm with six colors. So in Figure 2 we calculate the emission at the observer frequency  $\nu_{\text{obs},1} = 1.0 \times 10^{15}$  Hz and  $\nu_{\text{obs},2} = 5$  keV.

The synchrotron radiation of an i-shell interacting with the trail as a function of the observer time has been shown in Figure 2. What we observed is the emission comes from a series (several thousand) of i-shells interacting with the trail of slow neutrons rather than just from one. In the simplest model proposed here,  $\sim 3s(1+z)/\delta t = 300$  sample light curves overlap, each with a  $\delta t/(1+z) = 10^{-2}$  s delay. The resulting net observable fluxes at  $10^{15}$  Hz and 5 keV are about 0.11 Jy and  $1.4 \times 10^{-9}$  ergs  $\text{cm}^{-2} \text{s}^{-1}$  respectively. The signature of 5 keV emission is only marginal since the typical X-ray component during many GRBs is strong, too. So we do not discuss it any longer. The sensitivity of UVOT is about 19th magnitude in a 10-s exposure. The spacecraft's time-to-target is about 20–70 seconds, in principle, part of, if not all the emission predicted here can be detected in the near future. However, we need to investigate whether some other emissions during prompt  $\gamma$ -ray emission phase can dominate over the emissions predicted here or not. Recently, it is suggested that the relativistic  $e^\pm$  pair generated in the internal shocks phase

can power a UV flash  $\sim 13$ th magnitude, too (Fan & Wei 2004). But the actual value of that emission is proportional to  $\Gamma_m^4 [\delta t/(1+z)]^{5/2}$ , and in that work the fiducial value for these parameters are taken as 300 and  $0.1(1+z)$ s respectively. If  $\delta t \sim 0.01(1+z)$ s or smaller, as taken in this work, the UV flash predicted in Fan & Wei (2004) will be much weaker ( $\lesssim 18$ th magnitude) unless  $\Gamma_m$  is much larger than 300. Thus the UV emission predicted in this work can be detected independently.

### 3. DISCUSSION

In the standard baryonic internal shocks model for Gamma-ray bursts (Paczynski & Xu 1994; Rees & Mészáros 1994), the prompt  $\gamma$ -ray emission are powered by the interaction between ion shells with variable LFs. If these shells are neutron rich, the neutron component coupled with the "fast ions" is accelerated to a larger bulk LF ( $\sim$  several hundred), i.e., the fast neutron component, which is the focus of many publications (e.g., PD02, B03). However, the neutron component coupled with the "slow ions" can only be accelerated to a moderate bulk LF  $\sim$  tens, for which the typical  $\beta$ -decay radius  $\sim 10^{15}$  cm is significantly smaller than that of the fast neutrons. As long as the duration of GRB is long enough, the earlier n-shells will be swept by the much later i-shells successively till the radius  $\simeq R_{\beta,s}$ , where the slow neutrons have decayed significantly. Consequently the interaction may power observable emission.

In this Letter, with an extremely simplified model to describe the later but faster i-shells interacting with the much early but slow, decaying n-shells, we have shown that there comes a  $\sim 12$ th magnitude UV flash at the later time of long GRBs. That emission is bright enough to be detected by the upcoming Satellite SWIFT in the near future. The emission predicted here is independent on the poorly known medium distribution around the progenitor of GRBs, as long as it is not very dense; it is also independent of some other physical processes, such as the  $e^\pm$  pair emission (Fan & Wei 2004), the possible reverse shock emission (As usual, if we assume the number density of interstellar medium (ISM) is about  $1 \text{ cm}^{-3}$ ,  $R_{\beta,s}$  is much smaller than the deceleration radius  $\sim 10^{17}$  cm, at which about half of the kinetic energy of the ejecta has been converted into the thermal energy of the shocked ISM, for a neutron-free ejecta) and so on.

The resulting net light-curve in this work increases rapidly firstly. Then there comes a flat lasting  $\sim T_{90}$ , the duration of corresponding GRB. Finally, the light-curve drops sharply, where the observed emission are mainly contributed by the "Equal-arriving surface" effect (e.g., Kumar & Panaitescu 2000; Fan, Wei & Wang 2004), if other emission such as the reverse shock emission has not been taken into account. It should be noted that the model taken here is a great simplification of the real situation, in which both the ion/neutron shells are moving with variable LFs, it is natural to expect that the corresponding emission is variable with time, too. However, to simulate such complicate processes is far beyond the scope of this Letter. Nonetheless, the actual flux may not be far dimmer than our prediction since averagely speaking, the LFs of the slow n-shells and the i-shells are significantly

<sup>1</sup><http://swift.gsfc.nasa.gov/docs/swift/>

different, but their masses are comparable. So the interaction of them can power a bright UV flash in the range  $\sim 10^{13} - 10^{15}$  cm.

Y. Z. F thanks T. Lu and Z. Li for their long-term encouragement on the subject of neutron-fed GRBs. We also thank the anonymous referee for her/his constructive

comments. This work is supported by the National Natural Science Foundation (grants 10225314 and 10233010), the National 973 Project on Fundamental Researches of China (NKBRF G19990754). This work is in memory of Changlin, Zhang (Chang'an University, China), Y. Z. F's university physics teacher, who has passed away on 2004 September 6.

#### REFERENCES

- Bahcall, J. N., & Mészáros, P. 2000, *Phys. Rev. Lett.*, 85, 1362 (BM00)
- Beloborodov, A. M. 2003a, *ApJ*, 588, 931
- . 2003b, *ApJ*, 585, L19 (B03)
- Derishev, D. E., Kocharovsky, V. V., & Kocharovsky, V. L. V. 1999a, *A&A*, 345, L51
- . 1999b, *ApJ*, 521, 640
- Fan, Y. Z., & Wei, D. M. 2004, *MNRAS*, 351, 292
- Fan, Y. Z., Wei, D. M., & Wang, C. F. 2004, *A&A*, 424, 477
- Fuller, G. M., Pruet, J., & Abazajian, K. 2000, *Phys. Rev. Lett.*, 85, 2673
- Guetta, D., Spada, M., & Waxman, E. 2001, *ApJ*, 557, 399
- Kumar, P., & Panaitescu, A. 2000, *ApJ*, 541, L9
- Mészáros, P., & Rees, M. J. 2000, *ApJ*, 541, L5
- Paczyński, B., & Xu, G. H. 1994, *ApJ*, 427, 708
- Panaitescu, A., & Kumar, P. 2002, *ApJ*, 571, 779
- Piran, T. 1999, *Phys. Rep.*, 314, 575 (P99)
- Pruet, J., & Dalal, N. 2002, *ApJ*, 573, 770 (PD02)
- Rees, M. J., & Mészáros, P. 1994, *ApJ*, 430, L93
- Rossi, E. M., Beloborodov, A. M. & Rees, M. J. 2004, to appear in the 2003 Santa Fe GRB Conference Proceedings (astro-ph/0401355)
- Sari, R., & Esin, A. A. 2001, *ApJ*, 548, 787
- Wei, D. M., & Lu, T. 1998, *ApJ*, 505, 252
- . 2000, *A&A*, 360, L13
- Wijers, R. A. M. J., & Galama, T. J. 1999, *ApJ*, 523, 177
- Wu, X. F., Dai, Z. G., Huang, Y. F., & Lu, T. 2003, *MNRAS*, 342, 1131
- Zhang, B., & Mészáros, P. 2004, *Int. J. Mod. Phys. A.*, 19, 2385




FIG. 1.— The radiation frequencies involved in an i-shell interacting with the trail of slow n-shells as a function of observer time. The physical parameters are taken as:  $M_{\text{ion}} = 2M_{\text{n}}^0 = 5.6 \times 10^{26} \text{g}$ , which is corresponding to a Luminosity  $L = 10^{52} \text{ergs s}^{-1}$  and  $\delta t = 10^{-2}(1+z)\text{s}$ ,  $\Gamma_{\text{m}} = 200$ ;  $\Gamma_{\text{n,s}} = 30$ .  $\epsilon_{\text{e}} = 0.3$ ,  $\epsilon_{\text{B}} = 0.01$ ,  $z = 1$  ( $D_{\text{L}} = 2.2 \times 10^{28} \text{cm}$ ). The solid line, dashed line and dotted line represent  $\nu_{\text{c}}$  (the cooling frequency),  $\nu_{\text{a}}$  (the synchrotron self-absorption frequency) and  $\nu_{\text{m}}$  (the typical synchrotron frequency), respectively. The start point has been chosen to be  $R = 10^{14} \text{cm}$ , the corresponding start time  $t_{\text{obs,beg}} = 0.04(1+z)\text{s}$ . The end point has been chosen as  $2.5 \times 10^{15} \text{cm} \simeq 3R_{\beta,\text{s}}$ , the corresponding end time  $t_{\text{obs,end}} \simeq 3(1+z)\text{s}$ .




FIG. 2.— The synchrotron radiation powered by an i-shell interacting with the trail of slow n-shells as a function of observer time. The parameters are the same as those taken in Fig. 1. The solid line, dotted line represent  $\nu_{\text{obs}} = 10^{15}\text{Hz}$  and 5 keV respectively. For clarity, the value of dotted line has been multiplied by a factor 100. Please note that what we observed is the emission powered by a series (several thousand) of i-shells interacting with the slow neutrons' trail rather than just by one. In the simplest model proposed here, about 300 sample light curves overlap, each lags with  $\delta t = 10^{-2}(1+z)\text{s}$ . The resulting net observable fluxes at  $10^{15}\text{Hz}$  and 5keV are about 0.11Jy and  $1.4 \times 10^{-9}\text{ergs cm}^{-2} \text{ s}^{-1}$  respectively.